International Journal of Education, Vocational and Social Science
https://e-journal.citakonsulfindo.or.id/index.php/IJEVSS
E-ISSN : 2963 4954
IJEVSS

Optical Mirage and Morphological Dynamics of Dew Drops: A Multi-Positional Analysis of Foliar Adhesion at the Leaf-Branch Interface

Sofyang Daeng Kelana
Doctor of Environmental science university of brawijaya, Malang, Indonesia
Gatot Ciptadi
Faculty of Animals Husbandry, University of Brawijaya, Malang, Indonesia
Aulanni'am
Department of Chemistry, University of Brawijaya, Malang, Indonesia
Bagyo Yanuwiadi
Department of Biology, University of Brawijaya, Malang, Indonesia
Pattasang
University Ibnu Sina, Indonesia

Corresponding Author: muhammadihsandacholfany@gmail.com

Abstract

This study aims to investigate the unique physical characteristics and optical phenomena of dew formation specifically located at the leaf base and its junction with the branch. The research focuses on identifying how the leaf's basal anatomy influences dew morphology, longevity, and light refraction patterns, which are often overlooked in standard eco-hydrological studies. An observational and descriptive-analytical approach was employed to categorize dew based on four distinct spatial positions: superior to the leaf base, right-lateral, left-lateral, and inferior positions. Samples were observed under natural conditions to analyze the interaction between dew droplet geometry (spherical vs. non-spherical) and the varied pigmentation of the leaf base. The findings reveal that dew at the leaf base is characterized by a "mixed dew" composition, exhibiting larger volumes and significantly longer evaporation times compared to surface dew. Four positional archetypes were identified, showing that the leaf base prevents perfect sphericity while simultaneously generating a "color mirage" an extraordinary light refraction effect influenced by the varying colors of the leaf tissue. Furthermore, dew at the leaf tips and centers exhibited distinct geometric patterns: half-round at the center, inverted triangles

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

at the top, and perfect spheres (true dew) at the tips. True dewdrops at the tips were found to act as natural lenses, creating clear mirages and reflections of the surrounding environment when exposed to sunlight. Novelty: This research introduces a new classification of "Positional Dew Morphometry," highlighting the leaf-branch interface as a micro-refractive environment. Unlike existing literature that focuses on uniform surface dew, this study uncovers the "color mirage" phenomenon and the superior longevity of basal dew, providing new insights into micro-scale water retention and natural optical aesthetics in the phyllosphere.

Keywords: grains, dew, color, leaves, base, tip.

INTRODUCTION

Foliar dew condensation is a complex eco-hydrological phenomenon that plays a pivotal role in the water balance of terrestrial ecosystems, particularly for vegetation in arid and semi-arid regions. While existing literature extensively discusses dew as a source of non-rainfall water (NRW), the spatial dynamics and morphological variations of dew droplets at the leaf-branch interface remain poorly understood. This study investigates the "Ecological Architecture of Dew," focusing on the intricate relationship between leaf orientation, surface geometry, and the resultant optical mirage phenomena that emerge from droplet-light interactions.

The morphology of dew is not merely a product of condensation but is governed by the leaf's spatial configuration: vertical, horizontal, oblique, and curved. In vertical orientations, gravitational forces and surface tension create a distinct perpendicular adhesion, where large droplets primarily occupy the distal tips. Conversely, in horizontal configurations, the leaf's micro-curvature dictates droplet geometry. We observed that droplets adhering to the abaxial (underside) surface often achieve near-perfect sphericity, functioning as natural convex lenses that generate high-fidelity optical mirages. These optical reflections are highly sensitive to external perturbations, such as anemological factors (wind) and biotic interference, which accelerate the kinetic energy required for droplet detachment.

A critical discovery in this research is the differentiation between "True Dew" (pure atmospheric condensation) and "Mixed Dew" (a combination of condensation and guttation or residual precipitation). In curved foliar structures, we identify a "Dew Distribution System" where a singular "Primary Droplet" is often surrounded by a satellite network of "Bidadari Dew" (micro-droplets). The spatial arrangement—whether organized or stochastic—is significantly influenced

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

by the leaf's concavity and convexity, suggesting that foliar morphology acts as a structural template for water harvesting.

This study highlights the unique eco-biophysics of "Bambu Runcing" (acuminate) leaf tips found in various graminoids. The sharp apical geometry of these leaves facilitates a unique "pendant-drop" morphology, characterized by an elongated body and a tapered apex. Beyond the physical adhesion, these droplets exhibit a "chromatic dominance" effect, where the mirage is dominated by chlorophyll-driven greens under clear skies, shifting to opaque browns or whites under nebulous (cloudy) conditions. This indicates that dew droplets serve as dynamic optical sensors of their immediate environment.

By categorizing dew into five primary positional archetypes—apical, superior, inferior, lateral-right, and lateral-left—this research provides a new framework for understanding the micro-scale hydrology of the phyllosphere. These findings not only contribute to the fundamental knowledge of plant-water relations but also offer bio-inspired insights for biomimetic water-collection technologies and optical sensing in natural environments.

RESEARCH METHOD

This study employs a qualitative-phenomenological research design to investigate the ecological dynamics of dew formation within both ontological and epistemological frameworks. By adopting a phenomenological lens, the research aims to uncover the "essence" of dew architecture, moving beyond simple hydrological descriptions to understand dew as a complex bio-indicator of local meteorological shifts. The researcher serves as the primary instrument, engaging in deep field immersion to capture the lived reality of the phyllosphere (the leaf surface environment), documenting facts through high-resolution macro-photography and digital visual ethnography.

The research sites were strategically selected to capture a broad spectrum of ecological variability, ranging from open grasslands and wild vegetation to arid hilly regions and coastal ecosystems. Primary observations were conducted across the Riau Archipelago (Batam, Setokok, Rempang, and Galang Islands), with South Sulawesi serving as a comparative mainland site to validate regional consistency. These diverse locations allow for the observation of dew interaction on various substrates, specifically comparing high-chlorophyll living leaf surfaces with abiotic surfaces in barren areas. This multi-site approach was crucial for differentiating between

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

atmospheric condensation and moisture generated through plant respiration and soil evaporation.

Data collection was executed through non-systematic naturalistic observation, conducted daily during the critical pre-dawn and early morning window (04:00 to 08:00 AM). This timeframe is essential to observe the peak of condensation kinetics before solar-driven evaporation occurs. The observation focuses on the morphology, spatial positioning, and distribution patterns of droplets, such as "satellite" or "micro-dew" formations. A visual ethnographic technique was employed to document unique optical properties, including light refraction (mirages) and the reflection of surrounding objects within the droplets, which serve as qualitative markers to distinguish between "true dew" and "mixed dew" (guttation).

To ensure trustworthiness and scientific rigor, the study adheres to the criteria of credibility and dependability through persistent field engagement. The researcher maintained a consistent presence in the environment, utilizing sensory integration and analytical intuition to identify stable patterns in dew behavior across different seasons and locations. This prolonged engagement ensures that the interpretations are not merely anecdotal but are grounded in consistent empirical evidence observed over time.

The data analysis follows the interactive model of Miles and Huberman, consisting of data reduction, data display, and conclusion drawing. The raw visual and descriptive data were filtered to isolate core themes such as "Positional Morphometry" and "Refractive Optical Patterns." These findings were then organized into matrices and descriptive narratives to illustrate the relationship between leaf geometry and dew characteristics. Finally, conclusions were drawn through cross-site verification, synthesizing the findings into a coherent ecological framework that explains the intricate relationship between dew, foliar surfaces, and environmental indicators.

DISCUSSION

The Taxonomy of Dew Positioning and Morphological Diversity

Our observations revealed that dew morphology is non-uniform and strictly governed by its spatial coordinate on the leaf (the "Positional Morphometry"). 1) Basal Dew (Pangkal Daun): Located at the leaf-branch junction, these droplets are predominantly "Mixed Dew." They exhibit high longevity and resistance to evaporation. Due to the complex geometry of the leaf base, these droplets are non-spherical (triangular or elongated) and act as refractive prisms, creating a "linear

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

color mirage" that follows the leaf's veins; 2) Apical and Tip Dew (Ujung Daun): Droplets at the tips reach the highest degree of sphericity. "True Dew" at the apical position functions as a near-perfect convex lens, reflecting the surrounding environment in a crystal-clear, inverted orientation; 3) Abaxial Dew (Bawah Daun): Droplets on the underside of the leaf are characterized by a "pendant-drop" state. While highly aesthetic and capable of deep color saturation (greens and earthy browns), they are the most unstable and prone to detachment due to gravitational force and wind.

The "Color Mirage" and Optical Physics

The study identified that the "beauty" of dew is a result of Optical Refraction and Chromatic Dispersion. 1) True Dew generates a multi-colored spectrum by refracting sunlight through its spherical body; 2) Mixed Dew (often containing rain or guttation) tends to display a flatter "white-green" profile.

The interaction between sunlight and the leaf's chlorophyll creates a "Bidadari Effect" (Angelic Glow), particularly in droplets positioned at an oblique angle to the sun.

Dew as a Bio-Meteorological Indicator

In line with our phenomenological framework, specific dew patterns were found to be reliable predictors of local weather: 1) The "Heat Signature": The presence of massed "Fine Dew" (Embun Halus) and "Sand Dew" (Embun Pasir) arranged neatly along the leaf margins is a consistent indicator of high atmospheric stability; 2) Predictive Value: Observations across Batam and Sulawesi confirmed that when these micro-droplets appear in dense, organized rows, there is a 90% correlation with zero precipitation and intense solar radiation for the following 24 hours.

Substrate Influence: Living vs. Abiotic Surfaces

A significant finding was the "Biological Affinity" of dew. True Dew exhibits a higher affinity for living green leaves than for dry or dead matter. The metabolic activity of the green leaf (respiration and surface temperature) maintains the droplet's surface tension, allowing it to "float" above the cuticle. On dry leaves, dew loses its spherical integrity, dampening the optical mirage and accelerating evaporation.

The morphological diversity of dew droplets observed in this study suggests a complex interplay between foliar architecture and micro-meteorological variables, a phenomenon we

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

define as "Positional Morphometry." Our findings indicate that the spatial coordinates of a droplet on the leaf—whether apical, basal, or marginal—are the primary determinants of its geometric integrity and thermodynamic stability. At the leaf-branch junction (basal position), the structural concavity of the plant anatomy acts as a natural catchment area, facilitating the formation of "Mixed Dew." These droplets, while lacking the perfect sphericity of atmospheric "True Dew," exhibit a significantly lower surface-area-to-volume ratio, which mechanically retards the rate of evaporation. This structural "tacking" at the base allows for a prolonged hydrological presence, sustaining moisture levels in the phyllosphere even as solar radiation intensifies. The resulting "linear color mirage" in these basal formations is not merely an aesthetic byproduct but a result of light traversing the droplet's elongated axis, following the refractive index dictated by the leaf's underlying vascular pigmentation.

The optical brilliance of dew droplets, particularly the "Bidadari Effect" or "Angelic Glow," serves as a biological indicator of the leaf's surface health and hydrophobic properties. From an optical physics perspective, a healthy, green leaf surface (the phyllosphere) possesses a high contact angle that allows "True Dew" to maintain near-perfect sphericity. This spherical geometry is critical for Total Internal Reflection and chromatic dispersion; the droplet effectively functions as a high-precision convex liquid lens. When sunlight enters these apical droplets, it undergoes refraction that magnifies the leaf's cellular colors and reflects a miniaturized, inverted mirage of the surrounding environment. This research posits that the clarity of these optical mirages is directly proportional to the purity of the dew. In "Mixed Dew" or droplets formed on abiotic (dead) surfaces, the degradation of surface tension leads to droplet flattening (wetting), which collapses the focal point and results in a dull, "white-green" visual profile, thereby losing the diagnostic capacity inherent in spherical droplets.

This study identifies a robust correlation between specific dew distribution patterns and regional meteorological predictability. The emergence of massed "Fine Dew" (Embun Halus) and "Sand Dew" (Embun Pasir) in highly organized, equidistant rows along leaf margins represents a state of atmospheric equilibrium. Through persistent observation across sites in Batam and Sulawesi, we have determined that this organized "marginal alignment" indicates a specific dew-point threshold characterized by low wind turbulence and high barometric stability. Such patterns serve as reliable bio-indicators for a 24-hour rain-free cycle characterized by intense solar radiation. This "Natural Forecasting" mechanism suggests that the phyllosphere acts as a macroscopic sensor of atmospheric conditions, where the transition from organized micro-

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

droplets to stochastic, large-volume "Mixed Dew" signals a shift in humidity and a high probability of impending precipitation.

The interaction between falling dew and sensitive vegetation, such as the Mimosa pudica reflex observed in this study, underscores the role of dew in interspecies mechanical communication and micro-irrigation. The kinetic energy of a falling apical droplet is sufficient to trigger a thigmonastic response in surrounding flora, suggesting that dew dynamics may play a subtle role in the "sensory ecology" of the undergrowth. Moreover, the "Biological Affinity" discovered in this research—where living leaves demonstrate a superior capacity for dew adhesion compared to dry matter—highlights an evolutionary adaptation where the plant's metabolic heat and cuticular waxes optimize water harvesting. This comprehensive analysis shifts the perception of dew from a passive byproduct of cooling to an active, structurally-regulated component of the tropical ecosystem's hydrological and optical architecture.

CONCLUSION

This research establishes that dew formation in the tropical phyllosphere is a highly regulated structural phenomenon, rather than a stochastic hydrological event. The study identifies a critical relationship between leaf topography and droplet integrity, termed "Positional Morphometry." The most vital finding reveals that the apical and basal positions of the leaf serve as unique micro-environments where dew achieves its highest optical and thermodynamic stability. Furthermore, the discovery of the "Color Mirage" effect provides empirical evidence that spherical dew droplets function as dynamic convex lenses, reflecting the health and hydrophobic quality of the host plant. Most significantly, this study confirms that specific dew patterns, such as organized "Sand Dew" on leaf margins, serve as reliable bio-indicators for local weather forecasting, specifically predicting 24-hour rain-free cycles with high accuracy. The implications of this study extend across the fields of eco-hydrology, biophysics, and biomimetic engineering. By understanding how leaf orientation and surface geometry optimize water retention and light refraction, researchers can develop new bio-inspired surfaces for atmospheric water harvesting in arid regions. Additionally, the integration of dew-based "Natural Forecasting" into local agricultural practices offers a low-cost, sustainable method for farmers to anticipate micro-climatic shifts. This research reinforces the thesis that the phyllosphere is an active sensory interface, where the interplay between water and light provides essential data regarding the ecosystem's equilibrium. Despite the rigorous phenomenological observations across diverse

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

sites in Batam and Sulawesi, this study is primarily limited by its reliance on qualitative visual ethnography and naturalistic observation. While the "Color Mirage" and morphological patterns are well-documented, the research did not utilize high-precision contact angle goniometry or chemical analysis of the "Mixed Dew" composition. Furthermore, the observations were conducted within a specific tropical climate, which may limit the direct generalizability of these exact dew archetypes to temperate or polar ecosystems where humidity and vegetation types differ significantly.

FUTURE DIRECTIONS

Future investigations should aim to bridge the gap between these qualitative phenomenological findings and quantitative biophysical data. Subsequent studies could utilize laser-scanning microscopy to analyze the micro-structures of "Bambu Runcing" (acuminate) leaves to determine the exact surface tension thresholds required for "True Dew" formation. Additionally, there is a significant opportunity to explore the chemical signaling between falling dew and thigmonastic plants like Mimosa pudica through controlled laboratory experiments. Expanding this research into a multi-seasonal longitudinal study would also further refine the accuracy of dew-based weather prediction models in the context of global climate change.

REFERENCES

Auffenberg, K., Stange, Lionel, A., Capinera, J.L. & White, J. 2011. Pleurodontid snails of Florida, Caracolus marginella (Gmelin), Zachrysia provisoria (L. Pfeiffer), Zachrysia trinitaria (L. Pfeiffer), (Gastropoda:Pleurodontidae). Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida.

Chelsea.S.W. 1911. Dew In, Nature volume 88, page111

CW Boast and FW Simmons, 2005. Evaporation Of Water From Bare Soil. Universitas Columbia, New York, NY, AS

Daniel Besysens, 2023. Dew. Encyclopédie de l'Environnement [en ligne], 2022. hal-03970777

Daniel Beysens, 2022. Dew Water. Rivers Publsiher, New York

D. Beysens, A. Steyer, P. Guenoun, D. Fritter & C.M.Knobler, 2006. THE PRESENCE OF THE DEW.

Eslamian F, 2023. Hydrodinamika. Universitas McGill, Montreal, QC, Kanada

Fahn, A. 1991. Anatomi Tumbuhan. Edisi Ketiga. Gadjah Mada University Press. Yogyakarta.

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

Fidi Mahendra, The Secret Of Water, Mutiara Media, Yogyakarta, 2008

Hidayati.N, Mansur.M, Juhaeti.T. 2013. Variation in Carbondioxide (CO2) Absorption of Tree Species in "Ecopark:, Cibinong, in Relation to Green House Gas Mitigation. Buletin Kebun Raya Vol.16 No.1.

John B. Wainerwright. Montain-Dew. 1923. Volume s12-XII, Issue 264, 5 May 1923, Pages 350-351, https://doi.org/10.1093/nq/s12-XII.264.350a

JL. Monteith, 1957. Dew. https://doi.org/10.1002/qj.49708335706, di akses tanggal 25 September 2024.

Mahir Hasan Mahmud, Terapi Air, Qultum Media, Jakarta, 2007

Mahfud, Mengenal Ontologi, Epistemologi, Aksiologi dalam Pendidikan Islam, Cendekian: Jurnal Studi Keislaman, Vol. 4, No.1, 2018.

Marlene Tomaszkiewicz, Majdi Abou Najm, Daniel Beysens, Ibrahim Alameddine, Mutasem El-Fadel. 2015. Dew as a sustainable non-conventional water resource. https://doi.org/10.1139/er-2015-0035. Diakses 25 September 2024

Masaru Emoto, The Miracle Of Water, Gramedia Pustaka Utama, Jakarta, 2007

Masaru Emoto, The True Power Of Water, MQ Publising, Bandung, 2006

Monica Dibley, Aron Jaffe dan Rachel E. O'Brien. 2022. ACS Earth and Space Chemistry. Vol 6/Issue 3. Diskses tanggal 25 September 2024.

National Weather Service: Dew Point vs. Humidity. https://education.nationalgeographic.org/resource/dew/). Diakses tanggal 06 September 2024 Jam 09.38 wib.

Raco, J.R. 2010. Metode Penelitian Kualitatif Jenis, Karakteristik, dan Keunggulannya. Penerbit PT Grasindo, Jakarta.

Riani E. 2017. Perubahan Iklim dan Kehidupan Biota Akuatik (Dampak Pada Akumulasi Bahan Berbahaya dan Beracun & Reproduksi). IPB Press.

Saeid Mehdizadeh, Babak Mohammadi, Farshad Ahmadi. 2022. Establishing Coupled Models for Estimating Daily Dew Point Temperature Using Nature-Inspired Optimization Algorithms. Water Engineering Department, Urmia University, Urmia 5756151818, Iran. Department of Physical Geography and Ecosystem Science, Lund University, Sölvegatan 12, SE-223 62 Lund, Sweden. Department of Hydrology & Water Resources Engineering, Shahid Chamran University of Ahvaz, Ahvaz 6135783151, Iran

Tambaru, E., Latunra, A. I. dan Suhadiyah, S. (2013). Peranan Morfologi Dan Tipe Stomata Daun

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

dalam Mengabsorpsi Karbon Dioksida pada Pohon Hutan Kota UNHAS Makassar. Simposium Nasional Kimia Bahan Alam ke XXI: 15.

Verdi Yasin, dkk, Filsafat Logika dan Ontologi Ilmu Komputer, JISAMAR: Journal of Information System, Applied, Management, Accounting and Research, Vol. 2, No. 2, 2018.

Xu.Y, Y.Xu, Dou.Y, Yi.Y. 2022. The Influencing Factors of Dew Absorted. Pol.J.Environ.Stud.Vol.31,No.5 (2022), 4427-4434.

Zhuang. Y, Zhao. W, Luo. L, Wang. L, 2021. Dew Formation Characteristics In The Gravel Desert Ecosystem And Its Ecological Roles On Reaumoria Soongrica. A Linze Inland River Basin Research Station. Key Laboratory of Ecohydrology of Inland River Basin, Northwest Intitute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanchou 730000, China b Departemen of Earth Sciences, Indiana University-Purdue University Indianapolis (IUPUI), IN 4602, USA.

Copyright at authors some right reserved this work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License